

## SFB-Seminar

#### ZEIT:

30.11.2010, 16:00 Uhr - 19:00 Uhr

#### **ORT:**

HU Institut für Sportwissenschaft Philippstr. 13, 10115 Berlin (Mitte) Haus 11, Hörsaal 5

#### **PROGRAMM:**

#### 16:00 - 17:00 Marcos Jardim (IMECC-UNICAMP Brazil)

#### Moduli spaces of framed instantons sheaves on IPn

We generalize the ADHM construction of instantons to describe the moduli space of framed torsion-free

instanton sheaves on IPn, n ≥ 2. We then discuss instanton bundles on IP3 and describe the geometric

structure of their moduli space. If time allows, we shall also discuss perverse instanton sheaves.

This is joint work with Amar Henni and Renato Martins.

17:00 - 17:30 Kaffeepause/ Coffee Break

### 17:30 - 18:30 Giovanni Landi (Universita di Trieste)

# Dimensional Reduction over the quantum projective line and non-abelian q-vortices

We extend equivariant dimensional reduction techniques to the case of quantum spaces which are the

product of a Kähler manifold M with a quantum Riemann sphere. We work out the reduction of bundles

which are equivariant under the natural action of the quantum SU(2) group, and also of invariant gauge

connections on these bundles. The reduction of Yang-Mills gauge

#### Kontakt:

Humboldt-Universität zu Berlin . Institut für Mathematik SFB 647 . Unter den Linden 6 . 10099 Berlin Tel. +49 30 2093 1804 . Fax. +49 30 2093 2727 sfb647@math.hu-berlin.de theory on the product space leads to

q-deformations of the usual quiver gauge theories on M. In particular, we formulate generalized instanton

equations on the quantum space and show that they correspond to q-deformations of the usual

holomorphic chain vortex equations on M. We study some topological stability conditions for the

existence of solutions to these equations, and prove that the corresponding vacuum moduli spaces

are generally better behaved than their undeformed counterparts, but much more constrained by the

q-deformation. We work out explicit examples, including new examples of non-abelian vortices on

Riemann surfaces, and q-deformations of instantons whose moduli spaces admit the standard

hyper-Kähler quotient construction.